

Constraints on AdS/CFT Gravity Dual Models of Heavy Ion Collisions

Jorge Noronha¹, Miklos Gyulassy¹, and Giorgio Torrieri²

¹*Department of Physics, Columbia University, 538 West 120th Street, New York, NY 10027, USA*

²*Frankfurt Institute for Advanced Studies, Frankfurt am Main, Germany*

(Dated: June 22, 2009)

We show that the five-fold constraints due to (1) the observed nuclear modification of heavy quark jets measured via non-photonic electrons $R_{AA}^e(p_T \sim 6 \text{ GeV})$ in central Au+Au collisions at 200 AGeV, (2) the “perfect fluid” elliptic transverse flow of low transverse momenta pions, $v_2(p_T \sim 1 \text{ GeV})$ reported for noncentral collisions, (3) the global pion rapidity density dN_π/dy , (4) the lattice QCD entropy density deficiency, S/S_{SB} , of strongly coupled Quark-Gluon Plasmas (sQGP), and (5) a causal requirement are analytically correlated in a class of gauge/string gravity dual models of sQGP dynamics. Current RHIC/BNL and lattice QCD data are found to be remarkably compatible with these models if the t’Hooft and Gauss-Bonnet coupling parameters lie in the range $\lambda \approx 10 - 25$ and $0 < \lambda_{GB} < 0.09$. In addition, the observed five-fold correlation appears to favor color glass condensate over Glauber initial conditions within current systematic errors.

PACS numbers: 25.75.-q, 11.25.Tq, 13.87.-a

The combined observations of the quenching of hard (high transverse momentum or high quark mass jets) processes and the nearly “perfect fluid” elliptic flow of soft (low momentum transfer) hadrons produced in Au+Au collisions at $\sqrt{s} = 200 \text{ AGeV}$ at the Relativistic Heavy Ion Collider (RHIC) [1] have been interpreted as providing evidence for the formation of a new form of strongly interacting quark-gluon plasma (sQGP) [2]. Furthermore, bulk multiplicity (entropy) production systematics have been interpreted as evidence for gluon saturation of the initial conditions as predicted by the Color Glass Condensate (CGC) model [3, 4]. However, it has been a challenge to find a single consistent theoretical framework that can explain simultaneously both soft and hard phenomena. These phenomena include 1) the nuclear modification of high transverse momenta, $p_T > 5 \text{ GeV}$, jet observables, 2) the bulk collective flow observables ($p_T < 1 \text{ GeV}$), and 3) the sQGP thermodynamic equation of state entropy deficiency relative to the ideal Stefan-Boltzmann limit as predicted by nonperturbative Lattice QCD (LQCD). Attempts to explain these based on weak coupling (perturbative QCD) parton transport approaches [5, 6, 7, 8], especially the surprising small viscosity needed to fit the elliptic flow data, require large coupling extrapolations $\alpha_s = g_{YM}^2/4\pi \rightarrow 0.5 - 0.6$. At such large gauge couplings, on the other hand, the t’Hooft parameter $\lambda = g_{YM}^2 N_c \gg 1$ may already be large enough to validate string theory inspired AdS/CFT low energy approximations [9, 10]. The great theoretical advantage and appeal of gravity/gauge dual models [9]-[24] is that they have led (for the first time) to *analytic* connections between a wide variety of thermodynamic and nonequilibrium dynamic variables at strong coupling that were not yet realized with traditional gauge theory techniques.

In this Letter, we focus on predicted analytic connections between three fundamental properties of the sQGP: (1) its equation of state (entropy= S), (2) its

long wavelength transport coefficients (viscosity= η), and (3) coupling between long wavelength near-equilibrium “soft medium” properties and short-wavelength non-equilibrated “hard probes” (energy loss per unit length, dE/dx). We show that these three properties together with global entropy and causality restrictions can provide valuable phenomenological constraints of higher dimensional gravity dual models of sQGP in heavy ion collisions. We consider here the constraints imposed by current RHIC and LQCD data on a class of gravity dual models that include quadratic as well as quartic curvature corrections to the classical Einstein-Hilbert action for the effective 5 dimensional dual gravity action. Therefore, we implement and extend the suggestion made in [16] to include perturbatively both the lowest order $O(\lambda_{GB} \sim 1/N_c)$ Gauss-Bonnet $\mathcal{R}^2$ and $O(1/\lambda^{3/2})$ $\mathcal{R}^4$ curvature corrections to the three properties above.

Our analysis is based on the following remarkably simple algebraic expressions relating these three fundamental sQGP properties:

$$S/S_{SB} = \frac{3}{4} \left(1 + \lambda_{GB} + \frac{15}{8} \frac{\zeta(3)}{\lambda^{3/2}} \right), \quad (1)$$

$$\eta/s = \frac{1}{4\pi} \left(1 - 4\lambda_{GB} + 15 \frac{\zeta(3)}{\lambda^{3/2}} \right), \quad (2)$$

$$\tau_Q^{-1} = \mu_Q \left(1 + \frac{3}{2} \lambda_{GB} + \frac{15}{16} \frac{\zeta(3)}{\lambda^{3/2}} \right) \quad (3)$$

where $s = S/V$ is the entropy density. The heavy quark jet relaxation rate, $1/\tau_Q$, is controlled by $\mu_Q = \sqrt{\lambda} \pi T^2 / 2M_Q$ for a heavy quark with mass M_Q in a plasma of temperature T . The relaxation time is related to the energy loss per unit length through $\tau_Q(\lambda, \lambda_{GB}) = -1/(d \log p / dt) = -1/(d \log E / dx)$, where $p = M_Q \gamma v$ and $v = p/E$.

We note that the $\mathcal{R}^4$ correction $O(1/\lambda^{3/2})$ to the heavy quark jet energy loss is a new result [25] reported in this Letter and it is needed for a consistent additive perturba-




FIG. 1: (Color online) The gravity dual model correlation of the nuclear modification factor, R_{AA}^e , of hard ($p_T = 5.5$ GeV, 0-10% centrality) non-photonic electrons with the elliptic transverse flow moment, v_2 , of soft ($p_T = 1$ GeV, 20-60% centrality) pions assuming Glauber (left) and CGC (right) initial participant distributions and $dN_\pi/dy = 1000$ are compared. The green region corresponds to the range of entropy function deficiencies, S/S_{SB} , consistent with infinite volume extrapolations of lattice QCD data [34]. The red (blue) ellipse indicates PHENIX (STAR) 0-10% centrality R_{AA}^e and 20-60% v_2 centrality Au+Au 200 AGeV data from RHIC [35]-[38]. The dashed blue curves show fixed $S/S_{SB} = 0.8$ while for the solid black and dotted-dashed curves $S/S_{SB} = 0.85$ and $S/S_{SB} = 0.9$, respectively in the range $\lambda = 5, \dots, 30$. The causality bound curve with $\lambda_{GB} = 0.09$ is indicated by solid red. The gray ellipse shows preliminary v_2 for minimum bias data with estimated non-flow effects subtracted [39].

tive application of this class of AdS/CFT models to heavy ion reactions. We consider here only heavy quark observables because gravity dual string drag models [13] apply only for heavy $M_Q > \sqrt{\lambda}T$ jets. The heavy quark jet drag is then modeled as a trailing string moving in a black brane background according to the classical Nambu-Goto action (with dilaton neglected) $\mathcal{A}_{NG} = -\frac{1}{2\pi\alpha'} \int d^2\sigma \sqrt{-g}$ where $g = \det g_{ab} = G_{\mu\nu} \partial_a X^\mu \partial_b X^\nu$ is the induced worldsheet metric, $\sigma^a = (\tau, \sigma)$ are the internal worldsheet coordinates, $G_{\mu\nu}(X)$ is the background metric, and $X^\mu = X^\mu(\tau, \sigma)$ is the embedding of the string in spacetime. The trailing string ansatz (where $\tau = t$, $\sigma = u$ and $X^\mu(t, u) = (t, x_0 + vt + \xi(u), 0, 0, u)$) describes the asymptotic behavior of a string attached to a moving heavy quark (the string endpoint) with velocity v in the x direction and located at a fixed AdS radial coordinate $u_m \gg u_h$ [13]. The black brane horizon coordinate $u_h \propto T\alpha'$ is determined by $G_{00}(u_h) = 0$. Using the ansatz above and the string's classical equations of motion, one can show that the drag force $dp/dt = -Cv/(2\pi\alpha')$, where C is a constant determined by the negativity condition

that

$$g(u) = G_{uu} (G_{00} + v^2 G_{xx}) \left(1 + \frac{C^2 v^2}{G_{00} G_{xx}} \right)^{-1} < 0 \quad (4)$$

for $u_h \leq u \leq u_m$. However, both the numerator and denominator in Eq. (4) change their sign simultaneously at a certain u^* [13] given by the root of the equation $G_{00}(u^*) + v^2 G_{xx}(u^*) = 0$. This fixes $C = G_{xx}(u^*)$ and $dp/dt = -v G_{xx}(u^*)/(2\pi\alpha')$. Neglecting higher-order derivative corrections in $\mathcal{N} = 4$ SYM one finds $u^* = u_h \sqrt{\gamma}$, where $\gamma = 1/\sqrt{1-v^2}$. The condition that $u^* \leq u_m$ leads to a maximum “speed limit” for the heavy quark jet to be consistent with this trailing string ansatz given by $\gamma_{\max} \leq u_m^2/u_h^2$ [14].

Using the metric derived in [10] to $\mathcal{O}(\alpha'^3)$, one can compute the effects of quartic corrections on the drag force [15] and determine u^* perturbatively to $\mathcal{O}(\lambda^{-3/2})$ as [26] $u^* = u_h \sqrt{\gamma} \left[1 + \frac{15}{32} \frac{\zeta(3)}{\lambda^{3/2}} v^2 \left(5 + \frac{5}{\gamma^2} - \frac{3}{\gamma^4} \right) \right]$ and the drag force

$$\frac{dp}{dt} = -\sqrt{\lambda} T^2 \frac{\pi}{2} v \gamma \left[1 + \frac{15}{16} \frac{\zeta(3)}{\lambda^{3/2}} \left(1 - \frac{197}{24\gamma^4} + \frac{67}{24\gamma^6} \right) \right]. \quad (5)$$

The heavy quark mass at $T = 0$ is $M_Q = u_m/(2\pi\alpha')$ and, to leading order in $1/\lambda$, $u_m^2/u_h^2 \simeq \frac{4M_Q^2}{\lambda T^2}$. Thus, the corrected u^* displayed above defines a new speed limit $\gamma_m \simeq \frac{4M_Q^2}{\lambda T^2} \left[1 - \frac{5}{16} \left(\frac{4\pi\eta}{s} - 1 \right) \right]$, after neglecting terms of $\mathcal{O}(1/\gamma, 1/N_c)$. Note that γ_m and dp/dt decrease with increasing η/s .

For our applications, we consider the range $\lambda \sim 5 - 30$ and $|\lambda_{GB}| < 0.1$ with fixed $N_c = 3$. In this parameter range the $1/\lambda^{3/2}$ and $\lambda_{GB} \sim 1/N_c$ corrections are comparable. We neglect known (but formally) higher order terms [12] $\mathcal{O}(\sqrt{\lambda}/N_c^2)$ in this first attempt to test predicted dynamical correlations between hard and soft phenomena in high energy A+A collisions.

The small Gauss-Bonnet parameter $\lambda_{GB} = (c - a)/4c$ is related to the central charges c and a (related to the conformal anomaly in curved spacetime) of the dual CFT as noted in Eq. (2.14) of Ref. [16]. Varying λ_{GB} provides a parametric way to explore deformations of the original $\mathcal{N} = 4$ $SU(N_c)$ SYM theory. Interest in Gauss-Bonnet deformations were heightened when Kats and Petrov [21] argued that for $\mathcal{N} = 2$ $Sp(N_c)$, $\lambda_{GB} = 1/8N_c$, the KSS viscosity bound on $\eta/s \geq 1/4\pi$ was violated by 17% for $N_c = 3$. As further shown in [16], a large class of other effective CFTs are now known to lead to similar $\lambda_{GB} \propto 1/N_c$ effects. However, the analysis of Refs. [18]-[20] revealed that λ_{GB} deformations are limited by requirements of causality and positive energy flow to a narrow range $-7/36 < \lambda_{GB} < 9/100$.

In order to convert τ_Q into the observed nuclear modification of single non-photonic electrons, $R_{AA}^e(p_T = 5.5 \text{ GeV})$, from quenched heavy quark jets, we follow and extend [27] by using here the generalized drag force




FIG. 2: (Color online) Five fold phenomenological constraints in the t'Hooft and Gauss-Bonnet parameter space (λ, λ_{GB}) . (1) The green region with black dashed contours is from lattice QCD constraints on $0.8 < S/S_{SB} < 0.9$. (2) The cyan region is determined from noncentral elliptic flow $v_2(p_T = 1, 20 - 60\%) = 0.11 \pm 0.01$ [38]. Blue dashed contours correspond to fixed $4\pi\eta/s = 1, 1.5, 2$. The inversion $\lambda_{GB}(\lambda, v_2(\eta/s))$ is based on minimum bias viscous hydro results of [31] assuming CGC initial eccentricities scaled by a factor 1.1 in panel (a) and unscaled 1.0 in panel (b). The entropy is constrained by the (3) $dN_\pi/dy = 1000$ pion rapidity density in central collisions. (4) The gray region and contours are determined from central $R_{AA}^e(p_T = 5.5 \text{ GeV}) = 0.25 \pm 0.07$ data [35]. (5) The horizontal red line constraint is the causality upper bound for $\lambda_{GB}^{max} = 0.09$ [19, 20]. The yellow trapezoidal region with the purple boundary is the intersection of the five fold constraint bands. Note the red circled five fold conjunction area in panel (a) ($\lambda \approx 13, \lambda_{GB} \approx 0.08$) that is absent in the unscaled panel (b).

in Eq. (3) to compute the path length, L , dependent heavy quark fractional energy loss $\epsilon(L)$. The heavy quark jet nuclear modification factor is then $R_{AA} = \langle (1 - \epsilon)^{n_Q} \rangle_L$, where $n_Q(p_T)$ is the flavor dependent spectral index $n_Q + 1 = -\frac{d}{d \ln p_T} \ln \left(\frac{d\sigma_Q}{dy dp_T} \right)$ obtained from FONLL production cross sections [28] as used in [27]. The path length average of the nuclear modification at impact parameter b is computed using a Woods-Saxon nuclear density profile with Glauber profiles $T_A(\vec{x}_\perp)$ with $\sigma_{NN} = 42 \text{ mb}$. For 0-10% centrality triggered data both Glauber and CGC geometries lead to similar numerical results [29]. The distribution of initial hard jet production points at a given $\vec{x}_\perp$ and azimuthal direction ϕ is taken to be proportional to the binary parton collision density, $T_{AA}(\vec{x}_\perp, b)$. We assume a longi-

tudinally expanding local (participant) parton density $\rho(\vec{x}_\perp, b) = \chi \rho_{part}(\vec{x}_\perp, b)/\tau$, where $\chi \equiv (dN_\pi/dy)/N_{part}$ and ρ_{part} is the Glauber participant nucleon profile density. However, we evaluate (3) with a reduced temperature $T_{CFT} = 0.74(S/S_{SB})^{1/3} T_{QCD}$ to take into account the smaller number of degrees of freedom in a strongly-coupled QCD plasma, which is similar to the prescription given in [30]. We compute the heavy quark modification factor R_{AA}^Q via

$$R_{AA}^Q(p_T, b) = \int_0^{2\pi} d\phi \int d^2\vec{x}_\perp \frac{T_{AA}(\vec{x}_\perp, b)}{2\pi N_{bin}(b)} \times \exp \left[-n_Q(p_T) \int_{\tau_0}^{\tau_f} \frac{d\tau}{\tau_c(\vec{x}_\perp + \tau \hat{e}(\phi), \phi)} \right] \quad (6)$$

where N_{bin} is the number of binary collisions. Here, $\tau_0 = 1 \text{ fm/c}$ is the assumed plasma equilibration time and τ_f is determined from $T(\vec{\ell}, \tau_f) = T_f = 140 \text{ MeV}$, i.e, the time at which the local temperature falls below a freeze-out temperature taken from [31]. Systematic errors associated with the freeze-out condition will be discussed elsewhere.

In order to compute $v_2(1 \text{ GeV})$ for the $\mathcal{C}(20-60\%)$ centrality class, we employ a linear fit to the numerical viscous hydrodynamic results of Luzum and Romatschke [31]. The dependence of v_2 on viscosity for both Glauber [32] and CGC [4] initial transverse profiles can be well fit with

$$v_2(p_T, \eta/s, \mathcal{C}) = a(p_T) \epsilon_2(\mathcal{C}) (1 - b \eta/s) \quad (7)$$

where $\epsilon_2(\mathcal{C}) = \langle y^2 - x^2 \rangle_{\mathcal{C}} / \langle x^2 + y^2 \rangle_{\mathcal{C}}$ is the average initial elliptic geometric eccentricity for the centrality class $\mathcal{C}$. To rescale the minimum bias viscous hydro results of Ref. [31] to the considered 20-60% centrality class we use the factor $\epsilon_2^{Glaub}(20-60\%) / \epsilon_2^{Glaub}(0-92\%) = 0.317/0.281 = 1.128$ from Ref. [33]. Our fit to the rescaled numerical results of [31] give $b \approx 2.5$ and $a(p_T = 1) \epsilon_2(20-60\%) \approx 0.14$ (0.098) for CGC (Glauber) initial conditions. We consider this 20-60% centrality class because, as shown in Fig. 23 of [37], there is good agreement at $p_T \sim 1 \text{ GeV}$ between STAR $v_2(4)$ and PHENIX $v_2(BBC)$ data and non-flow effects [39] that complicate the interpretation of minimum bias data in [31] are reduced.

We find (see Figs. 1 and 2) that a (reasonable) combination of model parameters (λ and λ_{GB}) can account for the correlation between the reported R_{AA}^e [35, 36] and v_2 [38] taking into account the LQCD constraint on the equations of state deficiency S/S_{SB} [34] (in the range $T \sim 2 - 4T_c$). Our results suggest that within current systematic errors a small positive quadratic curvature correction with $0 < \lambda_{GB} < 0.1$ is preferred phenomenologically. Moreover, CGC initial eccentricities are favored over Glauber eccentricities for the centrality class considered. However, as emphasized by Fig. 2b, the results are rather sensitive to the $\lambda_{GB}(\lambda, v_2(\eta/s))$ inver-

sion. We find that a 10% reduction of the viscous hydrodynamic $v_2(\eta/s)$ (as for minimum bias centrality) virtually eliminates the yellow overlap region and could falsify the AdS/CFT description based on Eqs. (1-3). Great care is called for at this time to avoid premature conclusions. Systematic studies on viscous hydrodynamic dependence on centrality cuts and improved experimental control over non-flow corrections will be needed before definitive conclusions could be reached. The good news demonstrated by comparing Fig. 2a and 2b is that if improved theoretical and experimental control (better than 10%) over elliptic flow systematics and the nuclear modification of heavy quark jet observables can be reached, then rather strong experimental constraints on the AdS/CFT gravity dual model parameters could be achieved. We close by emphasizing [27] that future comparison of the nuclear modification of *identified* bottom and charm quark jets at RHIC and LHC combined with the fivefold (hard/soft) constraints considered in this Letter will provide especially stringent tests of AdS/CFT gravity dual phenomenology applied to high energy heavy ion reactions.

We thank A. Dumitru, S. Gubser, W. Horowitz, A. Poszkanzer, B. Cole, and W. Zajc for useful comments. J.N. and M.G. acknowledge support from US-DOE Nuclear Science Grant No. DE-FG02-93ER40764. G.T. acknowledges support from the Helmholtz International Center for FAIR within the framework of the LOEWE program (Landesoffensive zur Entwicklung Wissenschaftlich-Ökonomischer Exzellenz) launched by the State of Hesse.

-
- [1] I. Arsene *et al.* [BRAHMS Collaboration], Nucl. Phys. A **757**, 1 (2005); B. B. Back *et al.*, [PHOBOS Collaboration] Nucl. Phys. A **757**, 28 (2005); J. Adams *et al.* [STAR Collaboration], Nucl. Phys. A **757**, 102 (2005); K. Adcox *et al.* [PHENIX Collaboration], Nucl. Phys. A **757**, 184 (2005).
 - [2] M. Gyulassy and L. McLerran, Nucl. Phys. A **750**, 30 (2005); E. V. Shuryak, Nucl. Phys. A **750**, 64 (2005).
 - [3] D. Kharzeev, E. Levin and M. Nardi, Phys. Rev. C **71**, 054903 (2005); L. D. McLerran and R. Venugopalan, Phys. Rev. D **49**, 2233 (1994).
 - [4] A. Dumitru, E. Molnar, and Y. Nara, Phys. Rev. C **76**, 024910 (2007).
 - [5] P. Danielewicz and M. Gyulassy, Phys. Rev. D **31**, 53 (1985).
 - [6] D. Molnar and M. Gyulassy, Nucl. Phys. A **697**, 495 (2002) [Erratum-ibid. A **703**, 893 (2002)].
 - [7] A. Majumder, B. Muller and X. N. Wang, Phys. Rev. Lett. **99**, 192301 (2007).
 - [8] O. Fochler, Z. Xu and C. Greiner, arXiv:0806.1169 [hep-ph].
 - [9] J. M. Maldacena, Adv. Theor. Math. Phys. **2**, 231 (1998); E. Witten, Adv. Theor. Math. Phys. **2**, 253 (1998); S. S. Gubser, I. R. Klebanov and A. M. Polyakov, Phys. Lett. B **428**, 105 (1998).
 - [10] S. S. Gubser, I. R. Klebanov and A. A. Tseytlin, Nucl. Phys. B **534**, 202 (1998).
 - [11] A. Buchel, J. T. Liu and A. O. Starinets, Nucl. Phys. B **707**, 56 (2005); A. Buchel, Nucl. Phys. B **803**, 166 (2008).
 - [12] A. Buchel, Nucl. Phys. B **803**, 166 (2008); R. C. Myers, M. F. Paulos and A. Sinha, Phys. Rev. D **79**, 041901 (2009); A. Buchel, R. C. Myers, M. F. Paulos and A. Sinha, Phys. Lett. B **669**, 364 (2008).
 - [13] C. P. Herzog, A. Karch, P. Kovtun, C. Kozcaz and L. G. Yaffe, JHEP **0607**, 013 (2006); S. S. Gubser, Phys. Rev. D **74**, 126005 (2006); J. Casalderrey-Solana and D. Teaney, Phys. Rev. D **74**, 085012 (2006).
 - [14] J. Casalderrey-Solana and D. Teaney, JHEP **0704**, 039 (2007); S. S. Gubser, Nucl. Phys. B **790**, 175 (2008).
 - [15] J. F. Vazquez-Poritz, arXiv:0803.2890 [hep-th].
 - [16] A. Buchel, R. C. Myers and A. Sinha, JHEP **0903**, 084 (2009).
 - [17] A. Karch and E. Katz, JHEP **0206**, 043 (2002).
 - [18] M. Brigante, H. Liu, R. C. Myers, S. Shenker and S. Yaida, Phys. Rev. D **77**, 126006 (2008).
 - [19] M. Brigante, H. Liu, R. C. Myers, S. Shenker and S. Yaida, Phys. Rev. Lett. **100**, 191601 (2008).
 - [20] D. M. Hofman and J. Maldacena, JHEP **0805**, 012 (2008).
 - [21] Y. Kats and P. Petrov, arXiv:0712.0743 [hep-th].
 - [22] P. K. Kovtun, D. T. Son and A. O. Starinets, Phys. Rev. Lett. **94**, 111601 (2005).
 - [23] R. G. Cai, Phys. Rev. D **65**, 084014 (2002).
 - [24] K. B. Fadafan, JHEP **0812**, 051 (2008).
 - [25] J. Noronha *et al.*, to be published.
 - [26] The drag force in this case had been computed numerically in Ref. [15] but the analytical expression in Eq. (5) is, as far as we know, a new result.
 - [27] W. A. Horowitz and M. Gyulassy, Phys. Lett. B **666**, 320 (2008).
 - [28] M. Cacciari, P. Nason, R. Vogt, Phys. Rev. Lett. **95**, 122001 (2005); M. L. Mangano, P. Nason, G. Ridolfi, Nucl. Phys. B **373**, 295 (1992).
 - [29] A. Adil, H. J. Drescher, A. Dumitru, A. Hayashigaki and Y. Nara, Phys. Rev. C **74**, 044905 (2006).
 - [30] S. S. Gubser, Phys. Rev. D **76**, 126003 (2007).
 - [31] M. Luzum and P. Romatschke, Phys. Rev. C **78**, 034915 (2008); Erratum-ibid. **79**, 039903 (E) (2009).
 - [32] P. F. Kolb, U. W. Heinz, P. Huovinen, K. J. Eskola, and K. Tuominen, Nucl. Phys. A **696**, 197 (2001).
 - [33] K. Adcox *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **89**, 212301 (2002); D. d'Enterria, private communication.
 - [34] A. Bazavov *et al.*, arXiv:0903.4379 [hep-lat]; M. Cheng *et al.*, Phys. Rev. D **77**, 014511 (2008); C. Bernard *et al.*, Phys. Rev. D **75**, 094505 (2007); Y. Aoki, Z. Fodor, S. D. Katz and K. K. Szabo, JHEP **0601**, 089 (2006).
 - [35] S. S. Adler *et al.* [PHENIX], Phys. Rev. Lett. **96**, 032301 (2006); A. Adare *et al.* [PHENIX], Phys. Rev. Lett. **98**, 172301 (2007).
 - [36] B. I. Abelev *et al.* [STAR], Phys. Rev. Lett. **98**, 192301 (2007).
 - [37] A. Afanasiev *et al.* [PHENIX Collaboration], arXiv:0905.1070 [nucl-ex].
 - [38] B. I. Abelev *et al.* (STAR Collaboration), Phys. Rev. C **77**, 054901 (2008).
 - [39] A. Poszkanzer, private communication.